<b>Analyte Specifications:</b>	
IFN-γ	
(Option	1)


**Catalog Number:** 

92036


MultiCyt™ QBeads™ PlexScreen

Reconstituting the IFN- $\gamma$  standard in 2 mL of assay diluent results in a concentration of 5000 pg/mL. Standard curves for nowash and 1-wash protocols were performed in triplicate (+/- S.D.). The limit of detection (dashed line) is 3 standard deviations above the mean of samples without Standard (N=12). Results are typical and may vary based on specific assay conditions.

## **No-Wash Assay Performance**



## 1-Wash Assay Performance



Lowest Detectable Test Conc. (pg/mL): 4.9

For research purposes only. Not for resale.

Lowest Detectable Test Conc. (pg/mL): 4.9

Rev. 1 Jan 2014

Analyte Specifications: IFN-γ (Option 1)


**Catalog Number:** 

92036


MultiCyt™ QBeads™ PlexScreen

Reconstituting the IFN-γ standard in 2 mL of assay diluent results in a concentration of 5000 pg/mL. Standard curves for nowash and 1-wash protocols were performed in triplicate (+/- S.D.). The limit of detection (dashed line) is 3 standard deviations above the mean of samples without Standard (N=12). Results are typical and may vary based on specific assay conditions.

## **No-Wash Assay Performance**



1-Wash Assay Performance



Lowest Detectable Test Conc. (pg/mL): 4.9

Lowest Detectable Test Conc. (pg/mL): 4.9

For research purposes only. Not for resale.

Rev. 1 Jan 2014